

Take Home Suboxone (buprenorphine-naloxone)

Before you start, make sure you are very dope-sick (bad withdrawal)

Do you have at least 4 of these signs?

...or score yourself using the scale on Page 2



Joint and bone aches



Bad chills or sweating



Restless, can't sit still



Twitching, tremors or shaking



Big (enlarged) pupils



Goosebumps



Heavy yawning



Runny nose, tears in eyes



Feel sick, throwing up, or diarrhea

How long has it been?

It should be at least... 24 hours since you used heroin / fentanyl / oxycodone

48 hours since you used long-acting morphine (Kadian)

If you take methadone, talk to your Opioid Agonist Treatment (OAT) Clinic

Wait as long as you can! Start it too early and you might feel very, very sick

Once you are very dope-sick (in bad withdrawal)

How to take Suboxone:

- Put the tablet under your tongue
- Do not swallow it it won't work
- Do not take with alcohol or sleeping pills (sedatives)
- Do not take when sleepy

How soon it will work:

- Must melt under your tongue to work
- Takes up to 10 minutes to dissolve
- Takes 20 to 45 minutes to work
- **Always keep Suboxone away from children

	Day 2 and Day 3		
Step 1	Step 2	Step 3	
Place 1 tablet (2 mg*) under your tongue	Place 1 tablet (2 mg*) under your tongue	Still feel sick? Take 1 tablet every hour	Place <u>all</u> the tablets needed for Day 1 under your tongue
Wait 1 hour	Wait 1 hour	Stop when: You feel better or You have taken all 6 tablets (12 mg*)	at one time Can take up to 6 tablets each day
Step 1 + Step	Go to an OAT Clinic for more Suboxone (see Page 2)		

if you feel a lot worse at any time, stop taking Suboxone - Return to the Emergency Department

Subjective Opiate Withdrawal Scale (SOWS)*

Please score each of the statements according to how you feel right now on a scale of 1 to 4. Add up all your scores to get your total SOWS withdrawal score.

Scale: 0 = Not at all 1 = A little 2 = Moderately 3 = Quite a bit 4 = Extremely

	Time:					
Symptoms:		Score	Score	Score	Score	Score
6 2	I feel anxious					
	I feel like yawning					
	I am perspiring					
	My eyes are tearing					
	My nose is running					
	I have goosebumps					
	I am shaking					
*	I have hot flushes					
	I have cold flushes					
	My bones and muscles ache					
6-7	I feel restless					
	I feel nauseous					
	I feel like vomiting					
	My muscles twitch					
	I have stomach cramps					
	I feel like using now					
Му	SOWS score (total score):					

If your SOWS withdrawal score is **17 or more** \rightarrow You are ready to start, follow the instructions on Page 1. If your SOWS withdrawal score is **less than 17** \rightarrow Check your score again in 1 to 3 hours.

*Handelsman L et al. AM J Drug Alcohol Abuse.1987

Opioid Agonist Treatment (OAT) Clinics

opioid / igomot redunion (o/ri) omnos								
Abbotsford	604-743-0543	Langley	604-514-7940	New Westminster	604-527-2904			
203 - 31943 South Fraser Way		305-20300 Fraser Highway		330 East Columbia Street				
Burnaby	604-453-1930	Maple Ridge	604-467-5179	(Royal Columbian Wellness Centre)				
3935 Kincaid Street (Burnaby Hospital)		106 - 22838 Lougheed Highway		Surrey	604-580-4950			
Chilliwack	604-703-6976	Mission	604-814-5600	101-13670 94A Avenue (beside	Surrey Hospital)			
		3 rd Floor, 7298 Hurd Street (Mission Hospital)		White Rock	604-541-6844			
				15521 Russell Avenue, Russell Unit				

Rapid Access to Addictions Care (RAAC) Clinics

Fraser South 604-587-3755 (Opt. 3) | Fraser North 604-527-2904 | Vancouver 604-806-8867 | 3740 - 94A Avenue (beside Surrey Hospital) | 330 East Columbia Street (Royal Columbian Wellness Centre) | Vancouver 604-806-8867 | 2nd floor, 1081 Burrard Street (St. Paul's Hospital)

Any problems contacting your OAT Clinic? Have other Substance Use concerns? Call the **Substance Use Service Access Team** (SUSAT): **1-866-624-6478**

www.fraserhealth.ca

This information does not replace the advice given to you by your healthcare provider.