


Above Knee Amputation – Early Exercises

Abdominal Strengthening – Crunches

1. Lie on your back with arms crossed over your chest and knee bent.
2. Flatten your back by tightening stomach muscles and tilting your hips up.
3. Raise shoulders off floor, tucking chin to chest.
4. Hold 5 seconds.



Hip Extension – Leg Press Down

1. Lie on your back with a towel roll under your thigh and your other leg bent with the foot on the bed.
2. Press your thigh down into the towel while squeezing your buttocks together. As you get stronger, try to lift your buttocks off the bed.
3. Hold 5 seconds.




Hip Adduction – Leg Press In

1. Lie on your back with a pillow between your thighs.
2. Squeeze your thighs together and down.
3. Hold 5 seconds.



Hip Flexion – Leg Lift Up

1. Lie on your back. If you can, support yourself up on your elbows.
2. Bend one leg up towards your chest.
3. Lower the leg back down to the bed.
4. Repeat with the other leg.



Hip Abduction – Side Leg Lift


1. Lie on your side with your lower leg bent.
2. Lift your upper leg up. Keep your hip straight and your thigh turned slightly down.
3. Hold 5 seconds.



Above Knee Amputation – Early Exercises - *continued*


Hip Flexion-Extension - Side Lying

1. Lie on your side with your lower leg bent.
2. Bring your upper leg forward toward your chest.
3. Now reach it back as far as you can without arching your back.




Hip Flexor Stretch

1. Lie on your back with both legs bent up.
2. Grasp one thigh with both hands and lower the other thigh to rest flat on the bed. You will feel a stretch on the front of this leg.
3. Hold 5 seconds. Repeat with other leg.



Prone Lying


1. Roll onto your front, leading with your affected leg.
2. Keep your hips and thighs flat against the bed to stretch the front of your hips.
3. Try to stay on your front for at least 10 minutes 2 times daily.



Prone Lying – Hip Extension


1. Lie on your front.
2. Lift your thigh off bed. Hold 5 seconds. Return to start position.

Special Instructions: Do not arch low back



Chair Push Ups

1. Grip arm rests of chair.
2. Push up through arms to lift buttocks from chair seat.
3. Hold 5 seconds.



Images used and/or adapted with permission from Providence Health Care

www.fraserhealth.ca

This information does not replace the advice given to you by your healthcare provider.