

Dysphagia Minced Diet

| Name: | Date: | |
|------------------------------|-------|--|
| | | |
| Dietitian: | | |
| | | |
| Speech Language Pathologist: | | |
| | | |
| Occupational Therapist: | | |

What is Dysphagia?

Some people have trouble swallowing when eating solid food and/or drinking liquids. When you have trouble swallowing, this is called 'dysphagia' (sounds like dis-fay-jah).

Signs of dysphagia include coughing, choking, gurgly voice, throat clearing, runny nose, teary eyes, increased trouble breathing, unexplained rise in body temperature, the feeling of food being stuck in your throat when you eat, leftover food in your mouth after swallowing, and/or trouble starting your swallow. Please note, however, that not everyone who has dysphagia shows these signs.

When you have dysphagia, you might need to change what and how you eat to make swallowing easier and safer.

A swallowing problem could cause food or liquid to go into the lungs. This is called 'aspiration' (sounds like ass-per-ay-shun). If you aspirate, you could get a lung infection such as pneumonia (sounds like new-moan-yah).

Which food texture is safest for me?

We recommend you have a 'Dysphagia Minced' diet. This means all foods should be minced, mashed, very soft, or finely grated. Minced foods should not need much chewing.

| Other diet re | commendation | S. | | |
|---------------|--------------|----|--|--|
| | | | | |
| | | | | |
| | | | | |

What can I do to help make swallowing easier?

- Keep distractions to a minimum by eating in a calm and quiet place.
- Remember to swallow before talking.
- Sit upright while eating.
- Take small bites and sips.
- Eat and drink slowly.
- Swallow your mouthful completely before taking another bite or sip. You might need extra swallows for each bite or sip.
- Moisten foods by adding the following:
 - smooth gravy or sauce
 - butter, margarine, or oil
 - mayonnaise, sour cream, or yogurt

Foods you need to avoid

A. Foods of mixed consistencies:

These are foods that have both a solid and liquid component. Because of this, they are more difficult to swallow.

Some examples include:

- broth-based or cream-based soups with pieces of meat or vegetables

(Examples: chicken noodle, cream of mushroom)

- dry cereal with milk

(Examples: Rice Krispies®, Raisin Bran®)

- canned fruit in juice or syrup

(Example: fruit cocktail)

B. Sticky foods

These foods can stick to the roof of your mouth and are hard to push down with your tongue.

Some examples include:

- peanut butter, cream cheese, cheese spread
- toffee, caramel, jube-jubes, hard candy, chewing gum
- pizza, bagels
- dried fruit

C. Dry, crumbly, and crunchy foods:

These foods fall apart easily and can cause coughing and/or choking.

Some examples include:

- any bread or baked products that are dry, hard or contain hard and dry ingredients

(Examples: toast, cookies, crackers, scones, muffins, whole grain and multigrain bread, English muffins, croissants)

- dry cereals, including granola or muesli
- rice, couscous, quinoa
- popcorn, pretzels
- crackers, potato chips, taco chips or shells
- coconut, nuts, seeds
- raw vegetables

| Other foods to avoid: | | |
|------------------------|--|--|
| | | |
| Meal delivery options: | | |
| | | |
| | | |
| · | | |

Dysphagia Minced Foods

Note: If you have also been told to thicken fluids, please see the Thickened Fluids booklet for more on what drinks to choose.

| Food Group | Foods to Choose | Foods to Avoid |
|--------------------------|--|---|
| Milk and Alternatives | plain or smooth, flavoured yogurt plain or smooth pudding cottage cheese (drained) cheeses if used as an ingredient (e.g. grated hard cheese melted into a soup or casserole) | yogurt with nuts, seeds, or fruit all cheese including hard cheeses, soft cheeses, cheese spread, and cream cheese |
| Grain Products | cooked refined cereals (e.g. instant oatmeal, Cream of Wheat®, Cream of Rice®, Pablum® cereal) lightly toasted, crustless bread that has been buttered while warm soft pasta and noodles that are cut to the size of macaroni and with sauce rice with sauce or a part of a mixed dish naan or roti while warm or softened in sauce pureed congee crackers crumbled and soaked in soup plain, chopped rice noodle roll with sauce | cooked cereals or breads with bran, seeds, nuts, or fruits all dry cereals breads or pastries that are hard or crumbly (e.g. toast, scone, bagel, English muffin, croissant, Danish, cookies, cakes, Asian style buns or pastries) French toast, waffles dry or plain rice or pasta dry breads (e.g. pita bread, tortilla, naan, roti, chapatti) crackers, melba toast, rice cakes, chips, popcorn, pretzels sticky rice deep fried dumplings |

| Food Group | Foods to Choose | Foods to Avoid |
|--------------------------|--|--|
| Meat and Alternates | all meat and poultry that is moist, tender, and minced whole, tender boneless fish canned tuna and salmon minced or mashed with sauce soft, moist scrambled eggs mashed, hard cooked egg, well moistened with mayonnaise or sauce soft cooked, minced, moist legumes and lentils smooth hummus soft and medium tofu peanut butter or nut butter if used as an ingredient in cooking or baking strained rajma (kidney bean) or chana masala/cholay (chickpea) thick dahl (lentil curry) | tough, dry or fried meats bacon, sausage with skin battered, breaded, or fried meat, poultry, or fish all fish with bones hard cooked eggs except in minced egg salad nuts and seeds fried tofu peanut or nut butter unless used as an ingredient in cooking or baking |
| Fruits and Vegetables | applesauce and other fruit purées soft, canned fruit if minced, or mashed and drained ripe fresh banana ripe avocado stewed dried fruit if minced or mashed soft, well cooked vegetables if minced or mashed (e.g. broccoli, cauliflower, green beans, carrots, chopped spinach) mashed or boiled potatoes, pureed scalloped potatoes, skinless squash, yam | canned mixed fruit (e.g. fruit cocktail) fresh fruits with tough skins fresh juicy fruits (e.g. oranges, watermelon, grapes) all raw vegetables, including salads stringy, tough or fibrous vegetables (e.g. bok choi, gai lan, celery) vegetables with skins (e.g. corn, lima beans) fried potatoes, French fries fried vegetables |

| Food Group | Foods to Choose | Foods to Avoid |
|---------------------------------------|--|--|
| Soups, Mixed Dishes and Entrees | pureed or strained soups, congee, or kichari (lentil and rice soup) thickened to the fluid consistency recommended sandwiches with moist, minced, soft salad filling (e.g. fish or egg salad) soft, moist mixed dishes (e.g. casseroles) made with minced ingredients in smooth sauces or gravies soft frittatas and crustless quiches with allowed ingredients | all other soups that are not pureed or strained (e.g. chicken noodle, cream of mushroom) sliced meat or cheese sandwiches grilled sandwiches (e.g. grilled cheese) burgers, hot dogs stews, stir fries entrées with pastry unless soft and moistened with sauce (e.g. chicken pot pie) pizza |
| Desserts and Sweets | all soft desserts that will not crumble (e.g. puddings including tapioca pudding, rice pudding, custard, mousses) honey, jellies, sugar, syrups pie filling smooth, sweet soup desserts (e.g. black sesame or walnut soup) | desserts or baked goods with seeds, nuts, or dried fruits desserts that are dry, crumbly, or chewy (e.g. cakes, squares, bars, brownies, loaves, crisps, cobblers, pies, tarts, pastries, cookies, Rice Krispies® squares) hard pie crust or other pastry jam with seeds, marmalades |

If you have questions after leaving the hospital, Call 8-1-1 to speak to a Registered Dietitian.

For an interpreter, say you language in English. Wait until the interpreter comes on the phone.

Adapted with permission: Vancouver Coastal Health FI.710.M56 Dysphagia Minced Diet (April 2014)

www.fraserhealth.ca

This information does not replace the advice given to you by your healthcare provider.

Catalogue #265447 (December 2019) To order: patienteduc.fraserhealth.ca

